

Chapter 4

Declarative Interpretation

Outline

- Algebras (which provide a semantics of terms)
- Interpretations (which provide a semantics of programs)
- Soundness of SLD-resolution
- Completeness of SLD-resolution
- Least Herbrand models
- Computing least Herbrand models

What is an Interpretation?

```
direct(frankfurt, san_francisco).
```

```
direct(frankfurt, chicago).
```

```
direct(san_francisco, honolulu).
```

```
direct(honolulu, maui).
```

```
connection(X, Y) :- direct(X, Y).
```

```
connection(X, Y) :- direct(X, Z), connection(Z, Y).
```

$D = \{FRA, DRS, ORD, SFO, \dots\}$

$frankfurt_J = FRA, chicago_J = ORD, san\text{-}francisco_J = SFO, \dots$

$direct_I = \{(FRA, SFO), (FRA, ORD), \dots\}$

$connection_I = \{(FRA, SFO), (FRA, ORD), (FRA, HNL), \dots\}$

What is an Interpretation?

`add(X, 0, X) .`

`add(X, s(Y), s(Z)) :- add(X, Y, Z) .`

$D = \mathbb{IN}$

$0_J = 0$

$s_J : \mathbb{IN} \rightarrow \mathbb{IN}$ such that $s_J(n) = n + 1$

$\text{add}_J = \{(0, 0, 0), (1, 0, 1), (0, 1, 1), (1, 1, 2), \dots\}$

Another Example

`add(X, 0, X) .`

`add(X, s(Y), s(Z)) :- add(X, Y, Z) .`

$D = \{0, s(0), s(s(0)), \dots\}$

$0_J = 0$

$s_J : D \rightarrow D$ such that $s_J(t) = s(t)$

$\text{add}_J = \{(0, 0, 0), (s(0), 0, s(0)), (0, s(0), s(0)), (s(0), s(0), s(s(0))), \dots\}$

(This will be called a “Herbrand model”.)

Algebras

V set of variables, F ranked alphabet of function symbols: An **algebra** J for F (or pre-interpretation for F) consists of:

1. domain $:\Leftrightarrow$ non-empty set D
2. assignment of a mapping

$$f_J : D^n \rightarrow D$$

to every $f \in F^{(n)}$ with $n \geq 0$

State σ over D $:\Leftrightarrow$ mapping $\sigma : V \rightarrow D$

Extension of σ to $TU_{F,V}$ $:\Leftrightarrow$ $\sigma : TU_{F,V} \rightarrow D$ such that for every $f \in F^{(n)}$

$$\sigma(f(t_1, \dots, t_n)) = f_J(\sigma(t_1), \dots, \sigma(t_n))$$

Interpretations

F ranked alphabet of function symbols, Π ranked alphabet of predicate symbols:

An interpretation I for F and Π consists of:

1. algebra J for F (with domain D)
2. assignment of a relation

$$p_i \subseteq \underbrace{D \times \dots \times D}_n$$

to every $p \in \Pi^{(n)}$ with $n \geq 0$

Herbrand Universes and Bases

Recall $TU_{F,V} : \Leftrightarrow$ term universe over function symbols F , variables V

$TB_{\Pi,F,V} : \Leftrightarrow$ term base (i.e., all atoms) over predicate symbols Π and F , V

- **Herbrand universe** $HU_F : \Leftrightarrow TU_{F,\emptyset}$
- **Herbrand base** $HB_{\Pi,F} : \Leftrightarrow TB_{\Pi,F,\emptyset}$

Interpretations (Example)

Let P_{add} “add-program”.

$I_1, I_2, I_3, I_4, I_5,$ and I_6 are interpretations for $\{s, 0\}$ and $\{add\}$:

I_1 : $D_{I_1} = \mathbb{N}, 0_{I_1} = 0, s_{I_1}(n) = n + 1$ for each $n \in \mathbb{N}, add_{I_1} = \{(m, n, m + n) \mid m, n \in \mathbb{N}\}$

I_2 : $D_{I_2} = \mathbb{N}, 0_{I_2} = 0, s_{I_2}(n) = n + 1$ for each $n \in \mathbb{N}, add_{I_2} = \{(m, n, m * n) \mid m, n \in \mathbb{N}\}$

I_3 : $D_{I_3} = HU_{\{s, 0\}}, 0_{I_3} = 0, s_{I_3}(t) = s(t)$ for each $t \in HU_{\{s, 0\}},$
 $add_{I_3} = \{(s^m(0), s^n(0), s^{m+n}(0)) \mid m, n \in \mathbb{N}\}$

I_4 : $D_{I_4} = HU_{\{s, 0\}}, 0_{I_4} = 0, s_{I_4}(t) = s(t)$ for each $t \in HU_{\{s, 0\}}, add_{I_4} = \emptyset$

I_5 : $D_{I_5} = HU_{\{s, 0\}}, 0_{I_5} = 0, s_{I_5}(t) = s(t)$ for each $t \in HU_{\{s, 0\}}, add_{I_5} = (HU_{\{s, 0\}})^3$

I_6 : $D_{I_6} = \{0, 1\}, 0_{I_6} = 0, s_{I_6}(n) = n$ for each $n \in \{0, 1\}, add_{I_6} = \{(m, n, m) \mid m, n \in \{0, 1\}\}$

Logical Truth (I)

E **expression** $:\Leftrightarrow E$ atom, query, clause, or resultant

E expression, I interpretation, σ state:

E **true in I under σ** , written: $I \models_{\sigma} E$

$:\Leftrightarrow$

by case analysis on E :

- $I \models_{\sigma} p(t_1, \dots, t_n) :\Leftrightarrow (\sigma(t_1), \dots, \sigma(t_n)) \in p_I$
- $I \models_{\sigma} A_1, \dots, A_n :\Leftrightarrow I \models_{\sigma} A_i$ for every $i = 1, \dots, n$
- $I \models_{\sigma} A \leftarrow \underline{B} :\Leftrightarrow$ if $I \models_{\sigma} \underline{B}$ then $I \models_{\sigma} A$
- $I \models_{\sigma} \underline{A} \leftarrow \underline{B} :\Leftrightarrow$ if $I \models_{\sigma} \underline{B}$ then $I \models_{\sigma} \underline{A}$

Logical Truth (II)

E expression, I interpretation:

Let x_1, \dots, x_k be the variables occurring in E .

- $\forall x_1, \dots, \forall x_k E$ universal closure of E (abbreviated $\forall E$)
- $\exists x_1, \dots, \exists x_k E$ existential closure of E (abbreviated $\exists E$)
- $I \models \forall E \Leftrightarrow I \models_{\sigma} E$ for every state σ
- $I \models \exists E \Leftrightarrow I \models_{\sigma} E$ for some state σ
- E true in I (or: I model of E), written: $I \models E \Leftrightarrow I \models \forall E$

Logical Truth (III)

S , T sets of expressions, I interpretation:

- I **model** of S , written: $I \models S \Leftrightarrow I \models E$ for every $E \in S$
- T semantic (or: logical) **consequence** of S , written $S \models T \Leftrightarrow$ every model of S is a model of T

P program, Q_0 query, θ substitution:

- $\theta \upharpoonright_{\text{var}(Q_0)}$ **correct answer substitution** of $Q_0 \Leftrightarrow P \models Q_0\theta$
- $Q_0\theta$ **correct instance** of $Q_0 \Leftrightarrow P \models Q_0\theta$

Models (Example)

Let P_{add} “add-program” and let $I_1, I_2, I_3, I_4, I_5,$ and I_6 be the interpretations from slide 8.

- $I_1 \models P_{add}$ (since $I_1 \models_{\sigma} c$ for every clause $c \in P_{add}$ and state $\sigma : V \rightarrow \mathbb{N}$:
 - (i) $(\sigma(x), \sigma(0), \sigma(x)) \in add_{I_1}$ and
 - (ii) if $(\sigma(x), \sigma(y), \sigma(z)) \in add_{I_1}$ then $(\sigma(x), \sigma(y)+1, \sigma(z)+1) \in add_{I_1}$)
- $I_2 \not\models P_{add}$ (e.g. let $\sigma(x) = 1$, then $I_2 \not\models_{\sigma} add(x, 0, x)$
since $(\sigma(x), \sigma(0), \sigma(x)) = (1, 0, 1) \notin add_{I_2}$)
- $I_3 \models P_{add}$ (like for I_1 ; we call I_3 a (least) **Herbrand model**)
- $I_4 \not\models P_{add}$ (e.g. let $\sigma(x) = s(0)$, then $I_4 \not\models_{\sigma} add(x, 0, x)$
since $(\sigma(x), \sigma(0), \sigma(x)) = (s(0), 0, s(0)) \notin add_{I_4}$)
- $I_5 \models P_{add}$ (like for I_1 ; we call I_5 a Herbrand model)
- $I_6 \models P_{add}$ (like for I_1)

Semantic Consequences (Example)

Let P_{add} “add-program”.

- $P_{add} \models add(x, 0, x)$
(for every interpretation I : if $I \models P_{add}$ then $I \models add(x, 0, x)$, since $add(x, 0, x) \in P_{add}$)
- $P_{add} \models add(x, s(0), s(x))$
(for every interpretation I : if $I \models P_{add}$ then $I \models add(x, 0, x)$
and $I \models add(x, s(0), s(x)) \leftarrow add(x, 0, x)$ (instance of clause), thus $I \models add(x, s(0), s(x))$)
- $P_{add} \not\models add(0, x, x)$
(consider interpretation I_6 from slide 8 with $I_6 \models P_{add}$;
 $I_6 \not\models add(0, x, x)$, since e.g. $I_6 \not\models_{\sigma} add(0, x, x)$ for $\sigma(x) = 1$,
since $(\sigma(0), \sigma(x), \sigma(x)) = (0, 1, 1) \notin add_{I_6}$)

Towards Soundness of SLD-Resolution (I)

Lemma 4.3 (i)

Let $Q \xRightarrow[c]{\theta} Q'$ be an SLD-derivation step and $Q\theta \leftarrow Q'$ the resultant associated with it.

Then $c \vDash Q\theta \leftarrow Q'$

Proof.

Let $Q = \underline{A}, B, \underline{C}$ with selected atom B . Let $H \leftarrow \underline{B}$ be the input clause and $Q' = (\underline{A}, \underline{B}, \underline{C})\theta$.

Then

$c \vDash H \leftarrow \underline{B}$ (variant of c)

implies $c \vDash H\theta \leftarrow \underline{B}\theta$ (instance)

implies $c \vDash B\theta \leftarrow \underline{B}\theta$ (θ unifier)

implies $c \vDash (\underline{A}, B, \underline{C})\theta \leftarrow (\underline{A}, \underline{B}, \underline{C})\theta$ (“context” unchanged)

Towards Soundness of SLD-Resolution (II)

Lemma 4.3 (ii)

Let ξ be an SLD-derivation of $P \cup \{Q_0\}$. For $i \geq 0$ let R_i be the resultant of level i of ξ .

Then $P \vDash R_i$

Proof.

Let $\xi = Q_0 \xRightarrow{\theta_1} Q_1 \dots Q_n \xRightarrow{\theta_{n+1}} Q_{n+1} \dots$ Induction on $i \geq 0$:

$i = 0$: $R_0 = Q_0 \leftarrow Q_0 = \text{“true”}$, thus $P \vDash R_0$

$i = 1$: $R_1 = Q_0\theta_1 \leftarrow Q_1$; by Lemma 4.3 (i): $P \vDash R_1$

$i \rightsquigarrow i + 1$: $R_{i+1} = Q_0\theta_1 \dots \theta_{i+1} \leftarrow Q_{i+1}$ is a semantic consequence of resultant $Q_i\theta_{i+1} \leftarrow Q_{i+1}$ associated with $(i + 1)$ -st derivation step and $R_i\theta_{i+1} = Q_0\theta_1 \dots \theta_{i+1} \leftarrow Q_i\theta_{i+1}$, thus by Lemma 4.3 (i) and induction hypothesis: $P \vDash R_{i+1}$

Soundness of SLD-Resolution

Theorem 4.4

If there exists a successful SLD-derivation of $P \cup \{Q_0\}$ with $\text{CAS } \theta$, then $P \models Q_0\theta$.

Proof.

Let $\xi = Q_0 \xRightarrow{\theta_1} \dots \xRightarrow{\theta_n} \square$ be successful SLD-derivation.

Lemma 4.3 (ii) applied to the resultant of level n of ξ implies $P \models Q_0\theta_1 \dots \theta_n$ and

$$Q_0\theta_1 \dots \theta_n = Q_0(\theta_1 \dots \theta_n|_{\text{Var}(Q_0)}) = Q_0\theta.$$

Comparison to Intuitive Meaning of Queries

Corollary 4.5

If there exists a successful SLD-derivation of $P \cup \{Q_0\}$, then $P \models \exists Q_0$.

Proof.

Theorem 4.4 implies $P \models Q_0\theta$ for some CAS θ .

Then, $P \models Q_0\theta$

implies for every interpretation I : if $I \models P$, then $I \models Q_0\theta$

implies for every interpretation I : if $I \models P$, then $I \models \forall(Q_0\theta)$

implies for every interpretation I : if $I \models P$, then $I \models \exists Q_0$

implies $P \models \exists Q_0$

Towards Completeness of SLD-Resolution

To show completeness of SLD-resolution we need to syntactically characterize the set of semantically derivable queries.

The concepts of [term models](#) and [implication trees](#) serve this purpose.

Term Models

V set of variables, F function symbols, Π predicate symbols:

The **term algebra** J for F is defined as follows:

1. domain $D = TU_{F,V}$
2. mapping $f_J : (TU_{F,V})^n \rightarrow TU_{F,V}$ assigned to every $f \in F^{(n)}$ with
$$f_J(t_1, \dots, t_n) \Leftrightarrow f(t_1, \dots, t_n)$$

A **term interpretation** I for F and Π consists of:

1. term algebra for F
2. $I \subseteq TB_{\Pi,F,V}$ (set of atoms that are true; equivalent: assignment of a relation $p_I \subseteq (TU_{F,V})^n$ to every $p \in \Pi^{(n)}$)

I **term model** of a set S of expressions : \Leftrightarrow I term interpretation and model of S

Herbrand Models

The **Herbrand algebra** J for F is defined as follows:

1. domain $D = HU_F$
2. mapping $f_J : (HU_F)^n \rightarrow HU_F$ assigned to every $f \in F^{(n)}$ with
$$f_J(t_1, \dots, t_n) \Leftrightarrow f(t_1, \dots, t_n)$$

A **Herbrand interpretation** I for F and Π consists of:

1. Herbrand algebra for F
2. $I \subseteq HB_{\Pi, F}$ (set of ground atoms that are true)

I **Herbrand model** of a set S of expressions $:\Leftrightarrow I$ Herbrand interpretation and model of S

I **least Herbrand model** of a set S of expressions

$:\Leftrightarrow I$ Herbrand model of S and $I \subseteq I'$ for all Herbrand models I' of S

Implication Trees

implication tree w.r.t. program P

$:\Leftrightarrow$

- finite tree whose nodes are atoms
- if A is a node with the direct descendants B_1, \dots, B_n then $A \leftarrow B_1, \dots, B_n \in inst(P)$
- if A is a leaf, then $A \leftarrow \in inst(P)$

E expression, S set of expressions:

- $inst(E) :\Leftrightarrow$ set of all instances of E
- $inst(S) :\Leftrightarrow$ set of all instances of Elements $E \in S$
- $ground(E) :\Leftrightarrow$ set of all ground instances of E
- $ground(S) :\Leftrightarrow$ set of all ground instances of Elements $E \in S$

Implication Trees (Example)

Let P_{add} “add-program”, $n \in \mathbb{N}$, V set of variables, $t \in TU_{\{s,0\},V}$, and

$$\begin{array}{c} \mathcal{T} = \quad add(t, s^n(0), s^n(t)) \\ \quad \quad \quad | \\ \quad \quad \quad add(t, s^{n-1}(0), s^{n-1}(t)) \\ \quad \quad \quad \vdots \\ \quad \quad \quad \vdots \\ \quad \quad \quad add(t, s(0), s(t)) \\ \quad \quad \quad | \\ \quad \quad \quad add(t, 0, t) \end{array}$$

If $t \in HU_{\{s,0\}}$, then \mathcal{T} is **ground implication tree** w.r.t. P_{add} .

Implication Trees Constitute Term Model

Lemma 4.7

Consider term interpretation I , atom A , program P

- $I \models A$ iff $inst(A) \subseteq I$
- $I \models P$ iff for every $A \leftarrow B_1, \dots, B_n \in inst(P)$: if $\{B_1, \dots, B_n\} \subseteq I$ then $A \in I$

Lemma 4.12

The term interpretation

$\mathcal{C}(P) :\Leftrightarrow \{A \mid A \text{ is the root of some implication tree w.r.t. } P\}$ is a model of P .

Ground Implication Trees Constitute Herbrand Model

Lemma 4.26

Consider Herbrand interpretation I , atom A , program P

- $I \models A$ iff $\text{ground}(A) \subseteq I$
- $I \models P$ iff for every $A \leftarrow B_1, \dots, B_n \in \text{ground}(P)$, $\{B_1, \dots, B_n\} \subseteq I$ implies $A \in I$

Lemma 4.28

The Herbrand interpretation

$\mathcal{M}(P) := \{A \mid A \text{ is the root of some ground implication tree w.r.t. } P\}$ is a model of P .

Example

Let P_{add} “add-program”, and V set of variables.

The term interpretation

$$\begin{aligned} \mathcal{C}(P_{add}) &= \{add(t, s^n(0), s^n(t)) \mid n \in \mathbb{N}, t \in TU_{\{s,0\},V}\} \\ &= \{add(s^m(v), s^n(0), s^{n+m}(v)) \mid m, n \in \mathbb{N}, v \in V \cup \{0\}\} \end{aligned}$$

and the Herbrand interpretation

$$\begin{aligned} \mathcal{M}(P_{add}) &= \{add(t, s^n(0), s^n(t)) \mid n \in \mathbb{N}, t \in HU_{\{s,0\}}\} \\ &= \{add(s^m(0), s^n(0), s^{n+m}(0)) \mid m, n \in \mathbb{N}\} \end{aligned}$$

are models of P_{add} .

Correct Answer Substitutions versus Computed Answer Substitutions (Example)

Let P_{add} “add-program”, and $Q = add(u, s(0), s(u))$ query.

- $\theta = \{u/s^2(v)\}$ correct answer substitution of Q , since $P_{add} \models Q\theta = add(s^2(v), s(0), s^3(v))$ (in analogy to slide 13 with $x = s^2(v)$).
- SLD-derivation of $P_{add} \cup \{Q\}$:

$$add(u, s(0), s(u)) \xRightarrow{\theta_1} add(u, 0, u) \xRightarrow{\theta_2} \square$$
with $\theta_1 = \{x/u, y/0, z/u\}$ and $\theta_2 = \{x/u\}$,
thus $\eta = (\theta_1\theta_2)|_{\{u\}} = \epsilon$ is a computed answer substitution of Q .
- Thus, $Q\eta$ more general than $Q\theta$.
- In fact, no SLD-derivation of $P_{add} \cup \{Q\}$ can deliver correct answer substitution θ .

Completeness of SLD-Resolution for Implication Trees

Query Q is n -deep.

$:\Leftrightarrow$

every atom in Q is the root of an implication tree,
and n is the total number of nodes in these trees

Lemma 4.15

Suppose $Q\theta$ is n -deep for some $n \geq 0$. Then for every selection rule \mathcal{R} there exists a successful SLD-derivation of $P \cup \{Q\}$ with $\text{CAS } \eta$ such that $Q\eta$ is more general than $Q\theta$.

Completeness of SLD-Resolution (I)

Theorem 4.13

Suppose that θ is a correct answer substitution of Q . Then for every selection rule \mathcal{R} there exists a successful SLD-derivation of $P \cup \{Q\}$ with $\text{CAS } \eta$ such that Q_η is more general than $Q\theta$.

Proof. Let $Q = A_1, \dots, A_m$. Then: θ correct answer substitution of A_1, \dots, A_m

implies $P \vdash A_1\theta, \dots, A_m\theta$

implies for every interpretation I : if $I \vdash P$, then $I \vdash A_1\theta, \dots, A_m\theta$

implies $\mathcal{C}(P) \vdash A_1\theta, \dots, A_m\theta$ (since $\mathcal{C}(P) \vdash P$ by Lemma 4.12)

implies $\text{inst}(A_i\theta) \subseteq \mathcal{C}(P)$ for every $i = 1, \dots, m$ (by Lemma 4.7)

implies $A_i\theta \in \mathcal{C}(P)$ for every $i = 1, \dots, m$

implies $A_1\theta, \dots, A_m\theta$ is n -deep for some $n \geq 0$ (by def. of $\mathcal{C}(P)$)

implies claim (by Lemma 4.15)

Completeness of SLD-Resolution (II)

Corollary 4.16

Suppose $P \models \exists Q$.

Then there exists a successful SLD-derivation of $P \cup \{Q\}$.

Proof. $P \models \exists Q$

implies $P \models Q\theta$ for some substitution θ

implies θ correct answer substitution of Q

implies claim (by Theorem 4.13)

Least Herbrand Model

Theorem 4.29 $\mathcal{M}(P)$ is the least Herbrand model of P .

Proof. Let I be a Herbrand model of P and let $A \in \mathcal{M}(P)$.

We prove $A \in I$ by induction on the number i of nodes in the ground implication tree w.r.t. P with root A . Then $\mathcal{M}(P) \subseteq I$.

$i = 1$: A leaf implies $A \leftarrow \in \text{ground}(P)$
implies $I \not\models A$ (since $I \models P$)
implies $A \in I$

$i \rightsquigarrow i+1$: A has direct descendants B_1, \dots, B_n (roots of subtrees)

implies $A \leftarrow B_1, \dots, B_n \in \text{ground}(P)$ and $B_1, \dots, B_n \in I$ (induction hypothesis)

implies $A \leftarrow B_1, \dots, B_n \in \text{ground}(P)$ and $I \not\models B_1, \dots, B_n$

implies $I \not\models A$ (since $I \models P$)

implies $A \in I$

Ground Equivalence

Theorem 4.30 For every ground atom A : $P \models A$ iff $\mathcal{M}(P) \models A$.

Proof. “ \Rightarrow ”: $P \models A$ and $\mathcal{M}(P) \models P$ implies $\mathcal{M}(P) \models A$ (semantic consequence).

“ \Leftarrow ”: Show for every interpretation I : $I \models P$ implies $I \models A$.

Let $I_H = \{A \mid A \text{ ground atom and } I \models A\}$ Herbrand interpretation.

$I \models P$

implies $I \models A \leftarrow B_1, \dots, B_n$ for all $A \leftarrow B_1, \dots, B_n \in \text{ground}(P)$

implies if $I \models B_1, \dots, I \models B_n$ then $I \models A$ for all ...

implies if $B_1 \in I_H, \dots, B_n \in I_H$ then $A \in I_H$ for all ... (Def. I_H)

implies $I_H \models P$ (by Lemma 4.26; thus I_H Herbrand model)

implies $A \in I_H$ (since $A \in \mathcal{M}(P)$ and $\mathcal{M}(P)$ least Herbrand model)

implies $I \models A$ (by Def. I_H)

Complete Partial Orderings

Let $(\mathcal{A}, \sqsubseteq)$ be a partial ordering (cf. Slide 18 for Chapter 2).

- **a least element** of $X \subseteq \mathcal{A}$

$:\Leftrightarrow a \in X, a \sqsubseteq x$ for all $x \in X$

- **a least upper bound** of $X \subseteq \mathcal{A}$ (Notation: $a = \sqcup X$)

$:\Leftrightarrow a \in \mathcal{A}, x \sqsubseteq a$ for all $x \in X$ and a is the least element of \mathcal{A} with this property

$(\mathcal{A}, \sqsubseteq)$ **complete partial ordering** (CPO) $:\Leftrightarrow$

- \mathcal{A} contains a least element (denoted by \emptyset)
- for every increasing sequence $a_0 \sqsubseteq a_1 \sqsubseteq a_2 \dots$ of elements of \mathcal{A} , the set $X = \{a_0, a_1, a_2, \dots\}$ has a least upper bound

Some Properties of Operators

Let $(\mathcal{A}, \sqsubseteq)$ be a CPO.

operator $T: \mathcal{A} \rightarrow \mathcal{A}$ **monotonic**

$:\Leftrightarrow I \sqsubseteq J$ implies $T(I) \sqsubseteq T(J)$

operator $T: \mathcal{A} \rightarrow \mathcal{A}$ **finitary**

$:\Leftrightarrow$ for every infinite sequence $I_0 \sqsubseteq I_1 \sqsubseteq \dots$,

$$\bigsqcup_{n=0}^{\infty} T(I_n) \text{ exists} \quad \text{and} \quad T(\bigsqcup_{n=0}^{\infty} I_n) \sqsubseteq \bigsqcup_{n=0}^{\infty} T(I_n)$$

operator $T: \mathcal{A} \rightarrow \mathcal{A}$ **continuous** $:\Leftrightarrow T$ monotonic and finitary

I pre-fixpoint of $T : \Leftrightarrow T(I) \sqsubseteq I$

I fixpoint of $T : \Leftrightarrow T(I) = I$

Iterating Operators

Let $(\mathcal{A}, \sqsubseteq)$ be a CPO , $T: \mathcal{A} \rightarrow \mathcal{A}$, and $I \in \mathcal{A}$.

- $T \uparrow 0 (I) :\Leftrightarrow I$
- $T \uparrow (n + 1) (I) :\Leftrightarrow T(T \uparrow n (I))$
- $T \uparrow w (I) :\Leftrightarrow \sqcup_{n=0}^{\infty} T \uparrow n (I)$

$T \uparrow a :\Leftrightarrow T \uparrow a (\emptyset)$ (for $a = 0, 1, 2, \dots, w$)

By the definition of a CPO :

If the sequence $T \uparrow 0 (I), T \uparrow 1 (I), T \uparrow 2 (I), \dots$ is increasing, then $T \uparrow w (I)$ exists.

Theorem 4.22

If T is a continuous operator on a CPO , then $T \uparrow w$ exists and is the least prefixpoint of T and the least fixpoint of T .

Consequence Operator

Consider the CPO $(\{I \mid I \text{ Herbrand interpretation}\}, \subseteq)$.

Let P be a program and I a Herbrand interpretation. Then

$$T_P(I) := \{A \mid A \leftarrow B_1, \dots, B_n \in \text{ground}(P), \{B_1, \dots, B_n\} \subseteq I\}$$

Lemma 4.33

- (i) T_P is finitary.
- (ii) T_P is monotonic.

T_P -Characterization

Lemma 4.32

A Herbrand interpretation I is a model of P iff

$$T_P(I) \subseteq I$$

Proof.

$$I \models P$$

iff for every $A \leftarrow B_1, \dots, B_n \in \text{ground}(P)$:

$$\{B_1, \dots, B_n\} \subseteq I \text{ implies } A \in I \quad (\text{by Lemma 4.26})$$

iff for every ground atom A : $A \in T_P(I)$ implies $A \in I$

iff $T_P(I) \subseteq I$

Characterization Theorem

Theorem 4.34

- $\mathcal{M}(P)$ (i)
- = least Herbrand model of P (ii)
- = least pre-fixpoint of T_P (iii)
- = least fixpoint of T_P (iv)
- = $T_P \uparrow \omega$ (v)
- = $\{A \mid A \text{ ground atom, } P \models A\}$ (vi)

Success Sets

success set of a program P : \Leftrightarrow

$\{A \mid A \text{ ground atom, } \exists \text{ successful SLD-derivation of } P \cup \{A\} \}$

Theorem 4.37

For a ground atom A , the following are equivalent:

- (i) $\mathcal{M}(P) \models A$
- (ii) $P \models A$
- (iii) Every SLD-tree for $P \cup \{A\}$ is successful
- (iv) A is in the success set of P

Objectives

- Algebras (which provide a semantics of terms)
- Interpretations (which provide a semantics of programs)
- Soundness of SLD-resolution
- Completeness of SLD-resolution
- Least Herbrand models
- Computing least Herbrand models